

**DACONIL WEATHERSTIK**

Product registration number: *MAPP 12636*  
*PCS 02680*



A suspension concentrate formulation containing 500 g/l (40.2% w/w) chlorothalonil

A broad spectrum fungicide for the control of many diseases in turf.

*The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work (UK only).*

This product label is compliant with the CPA Voluntary Initiative (VI) guidance. (UK only)



Net contents

Approval Holder and Marketing Company (UK)	Marketing Company (Ireland)
Syngenta Crop Protection UK Limited CPC4, Capital Park, Fulbourn, Cambridge, CB21 5XE Tel: Cambridge (01223) 883400	Syngenta Ireland Ltd Block 6, Cleaboy Business Park, Old Kilmeaden Road, Waterford, Ireland Tel: 051 377203

In case of toxic or transport emergency ring +44 (0)1484 538444 anytime

**PROTECT FROM FROST**  
**SHAKE WELL BEFORE USE**

Product code / Print date / xxxxxx

Batch Number

## **SAFETY PRECAUTIONS**

(references to COSHH and LERAP apply to UK use only)

### **(a) Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when applying by vehicle mounted/drawn equipment.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND RUBBER BOOTS when using a pedestrian controlled sprayer.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), AND SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH ALL PROTECTIVE CLOTHING thoroughly after use especially the insides of gloves.

WASH SPLASHES from skin or eyes immediately.


DO NOT BREATHE SPRAY.

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking and after work.

### **(b) Environmental protection**

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.



DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) Scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for inspection for three years.

**(c) Storage and disposal**

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add the washings to the sprayer tank at the time of filling and dispose of container safely.

DO NOT RE-USE CONTAINER for any other purpose

## DACONIL WEATHERSTIK



### **HARMFUL DANGEROUS FOR THE ENVIRONMENT**

A suspension concentrate formulation containing 500 g/l (40.2% w/w) chlorothalonil

**HARMFUL BY INHALATION.  
IRRITATING TO EYES AND RESPIRATORY SYSTEM  
MAY CAUSE SENSITISATION BY SKIN CONTACT  
LIMITED EVIDENCE OF CARCINOGENIC EFFECT  
VERY TOXIC TO THE AQUATIC ORGANISMS, MAY CAUSE LONG-TERM  
ADVERSE EFFECTS IN THE AQUATIC ENVIRONMENT**

Keep out of reach of children

When using do not eat, drink or smoke

Keep away from food, drink and animal feeding stuffs

Use appropriate containment to avoid environmental contamination.

Wear suitable protective clothing and gloves.

This material and its container must be disposed of in a safe way.

**To avoid risks to man and the environment, comply with the instructions for use.**

**MAPP 12636  
PCS 02680**

## **IMPORTANT INFORMATION**

### **FOR USE ONLY AS A HORTICULTURAL FUNGICIDE**

For use on: Turf

Maximum individual dose: 300 ml product per 100 sq. m.

Other specific restrictions: This product must only be applied by pedestrian controlled sprayers or vehicle mounted/drawn equipment. Do not apply with a knapsack sprayer.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

This leaflet is part of the approved Product Label.

## **DIRECTIONS FOR USE**

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

### **RESTRICTIONS**

DACONIL WEATHERSTIK is only for use on established turf.

DACONIL WEATHERSTIK must only be applied using standard vehicle-mounted spray equipment or pedestrian controlled sprayers. Do not apply with a knapsack sprayer.

DO NOT MOW within 24 hours after treatment.

Do not irrigate for 24 hours after treatment or until the spray has thoroughly dried on the grass leaves.

Do not tank mix with other pesticide sprays, surfactants or liquid fertilizers.

### **AREAS OF USE**

DACONIL WEATHERSTIK controls many major fungal diseases of fine turf such as golf greens, golf tees, bowling greens, cricket squares and ornamental lawns. It is also suitable for use on extensive grass such as golf fairways and sports pitches.

DACONIL WEATHERSTIK is only recommended for treatment on established turf.

### **DISEASES CONTROLLED**

DACONIL WEATHERSTIK is a broad spectrum fungicide for the control of many major diseases affecting grasses and especially those attacking intensively managed ornamental or amenity turf.

Being primarily a contact fungicide DACONIL WEATHERSTIK is best applied as a preventative treatment when the disease is at a very early stage of development. Where possible diseases should be positively identified. A knowledge of previous disease occurrence will also aid decisions regarding fungicide treatment. Although DACONIL WEATHERSTIK can be applied at any time of year, spring and autumn are usually when conditions are most favourable for disease development and when treatment is most appropriate.

### **APPLICATION RATE AND DISEASES CONTROLLED**

#### **Preventative treatment**

Application at first signs of disease: Apply 300 ml of DACONIL WEATHERSTIK in a minimum of 5 litres of water per 100 sq. m. Repeat at 3 – 6 week intervals if re-infection occurs, for the control of: Fusarium Patch or Pink Snow Mould (*Microdochium nivale*), Dollar Spot (*Sclerotinia homoeocarpa*), Red Thread (*Laetisaria fuciformis*), and Anthracnose (*Colletotrichum graminicola*).

### **Curative Treatment**

Application to established disease: Apply 300 ml of DACONIL WEATHERSTIK in a minimum of 5 litres water per 100 sq. m. Repeat at 1 – 2 week intervals as necessary for the control of Dollar Spot (*Sclerotinia homoeocarpa*), and for the reduction of Red Thread (*Laetisaria fuciformis*).

Note: DACONIL WEATHERSTIK is not recommended for the curative treatment of Fusarium Patch or Pink Snow Mould (*Microdochium nivale*), and Anthracnose (*Colletotrichum graminicola*).

### **MIXING**

Shake container thoroughly before use. Half fill spray tank with clean water, add the required quantity of DACONIL WEATHERSTIK and agitate to ensure complete mixing.

### **OTHER INFORMATION**

1. Some diseases can quickly damage turf. Treatment at a late stage of disease development will be more difficult and can leave bare soil patches needing renovation.
2. Use preventative sprays, especially against diseases which occur in winter and early spring.
3. If diseases recur regularly, check management practices, especially fertilizer treatment as this can affect disease occurrence if either in excess or deficient.